

On a Class of Singular Boundary Value Problems Which Contains the Boundary Conditions $X'(0) = X(1) = 0^*$

ANTONIO TINEO

*Universidad de los Andes, Facultad de Ciencias,
Departamento de Matematicas, Merida 5101, Venezuela*

Received October 29, 1990; revised September 9, 1992

0. INTRODUCTION

In this paper we prove an existence theorem for the boundary value problem

$$x'' + f(t, x) = 0 \quad (0.1)$$

$$x(0) = 0, \quad \lambda x(1) + \mu x'(1) = 0, \quad (0.2)$$

where $\lambda, \mu \geq 0$, $\lambda + \mu > 0$ and $f: (0, 1) \times (0, \infty) \rightarrow (0, \infty)$ is a continuous function which does not have, in general, a continuous extension to $[0, 1] \times [0, \infty)$. In this case, we say that problem (0.1) is singular.

By the change of variables $y(t) = x(1 - t)$, we get a parallel result for the boundary conditions

$$\lambda x(0) - \mu x'(0) = 0, \quad x(1) = 0, \quad (0.3)$$

and we apply this to proving some existence theorems for the following boundary value problem:

$$x'' + ct^{-1}x' + f(t, x) = 0, \quad x'(0) = x(1) = 0. \quad (0.4)$$

See Theorems 0.4–0.5 below.

These problems appear in a variety of applications. See [2, 5, 6], where $f(t, x)$ has the form $\alpha(t)x^{-p}$, for some $p > 0$ and some continuous function $\alpha: (0, 1) \rightarrow (0, \infty)$. See also [1, 3, 4, 7] for more general f . Problem (0.4) also arises in the search for positive radially symmetric solutions to the problem $[\Delta u + f(|x|, u) = 0, x \in \Omega, u|_{\Gamma} = 0]$, where Ω is the open unit ball

* This work was sponsored by the C.D.C.H.T. of Univ. de los Andes, Merida, Venezuela.

in \mathbb{R}^{c+1} (if c is an integer) centered at the origin, Γ is its boundary, and $|x|$ is the Euclidean norm of x . See [3, 4].

In the following we write $E_1 = C^2(0, 1) \cap C^1(0, 1] \cap [0, 1]$ and $E_0 = C^2(0, 1) \cap C^1[0, 1] \cap C[0, 1]$. We shall prove:

0.1. THEOREM. *Suppose that there are continuous functions $\alpha, \gamma: (0, 1) \rightarrow (0, \infty)$ and a decreasing function $g: (0, M) \rightarrow (0, \infty)$ (some $0 < M \leq \infty$); such that*

$$\gamma(t) \leq f(t, x) \leq \alpha(t) g(x) \quad \text{in } (0, 1) \times (0, M). \quad (0.5)$$

Assume further that $t\alpha(t)$ is integrable over $[0, 1]$ and

$$\int_0^M dx/g(x) > \int_0^1 t\alpha(t) dt. \quad (0.6)$$

If $\mu > 0$, then (0.1)–(0.2) has a solution u in E_1 such that $u(t) \leq M$ in $[0, 1]$.

Remark. If $M = \infty$ in (0.5), then (0.6) is automatically true. Moreover, in this case there is $\infty > N > 0$ such that (0.6) is true if we replace M by N . So, we can assume that M is finite.

From the change of variables $y(t) = x(1 - t)$, we get:

0.2. THEOREM. *Assume (0.5) and $\mu > 0$. If $(1 - t)\alpha(t)$ is integrable over $[0, 1]$ and*

$$\int_0^M dx/g(x) > \int_0^1 (1 - t)\alpha(t) dt, \quad (0.7)$$

then (0.1), (0.3) has a solution u in E_0 such that $u(t) \leq M$.

0.3. Remarks. (a) If $\mu = 0$ we can prove that (0.1)–(0.2) has one solution in $C^2(0, 1) \cap C[0, 1]$ if $t(1 - t)\alpha(t)$ is integrable over $[0, 1]$. This result appears in [7].

(b) Assume that $\alpha(t)g(kt)$ is integrable over $[0, 1]$ for all $k > 0$. Then, $t\alpha(t)$ is integrable over $[0, 1]$ and each solution u of (0.1)–(0.2) belongs to $C^1[0, 1]$. (Compare with Theorem 2.2 of [3].)

Proof. Since $g(x)/x$ is decreasing in $(0, \infty)$ then $g(x) \geq g(1)x$ in $(0, 1)$ and thus, $t\alpha(t)$ is integrable over $[0, 1]$. On the other hand, it is easy to prove that there exists $k > 0$, such that $u(t) \geq kt$, in $[0, 1]$. From this,

$$|u'(t) - u'(s)| \leq \int_s^t \alpha(\xi) g(k\xi) d\xi$$

for $0 < s < t < 1$, and then u' is uniformly continuous on $(0, 1)$. Consequently, u' has a continuous extension to $[0, 1]$, and the proof follows easily.

We have a parallel remark for Theorem 0.2.

(c) Let $f(t, x) = [x^2 + (a - t)^2] x^{-3}$, where $a = (2\lambda + \mu)^{-1} (\lambda + \mu)$. Then, $f(t, x)$ is decreasing in $x > 0$ and $f(t, x) \leq K(x^{-1} + x^{-3})$ for some constant $K > 0$. Thus, f satisfies the hypotheses of theorem 0.1. In this case, the unique solution to (0.1)–(0.2) is given by $u(t) = (2at - t^2)^{1/2}$. (See Corollary 2.3 below.) Note that $u \notin C^1[0, 1]$.

EXAMPLE. Let $f(t, x) = \alpha(t) \phi(x) x^{-p}$; where $p > 0$ and $\alpha: (0, 1) \rightarrow (0, \infty)$, $\phi: [0, \infty) \rightarrow (0, \infty)$ are continuous functions such that $t\alpha(t)$ is integrable over $[0, 1]$ and ϕ is increasing. Further, if $\phi(x) \rightarrow \infty$ as $x \rightarrow \infty$, assume that ϕ is differentiable on $(0, \infty)$ and $\phi'(x)/x^p \rightarrow 0$ as $x \rightarrow \infty$. Then, $f(t, x)$ satisfies the hypotheses in theorem 0.1. (This includes the "model case", $\phi \equiv 1$.)

Proof. Let I denote the integral of $t\alpha(t)$ over $[0, 1]$, and let us consider the following cases:

(1) ϕ is bounded. In this case let us choose $K, M > 0$ such that $\phi(x) \leq K$ for $x > 0$ and

$$M^{p+1} > (p+1)KI. \quad (0.8)$$

(2) ϕ is unbounded. In this case, by L'Hopital's rule,

$$\lim_{x \rightarrow \infty} (p+1)\phi(x)/x^{p+1} = \lim_{x \rightarrow \infty} \phi'(x)/x^p = 0,$$

and so there exists $M > 0$ such that

$$(p+1)\phi(M)I < M^{p+1}. \quad (0.9)$$

Now, let us define $g(x) = \phi(M)x^{-p}$ for $0 < x < M$ and $\gamma(t) = \phi(0)\alpha(t)/M^p$. Then, g is decreasing; (0.5) is satisfied and (0.6) is equivalent to (0.8) (resp. (0.9)) if ϕ is bounded (resp. unbounded). So, the proof is complete.

Theorem 0.1 is related to some results in [1, 3, 7]. In fact, our proof take some ideas from [1, 7].

Using Theorem 0.2 and a comparison result (Theorem 2.1 below), we shall prove the following:

0.4. THEOREM. Let $C \in C^0[0, 1] \cap C^1(0, 1)$ be an increasing function, which is positive in $(0, 1]$. If the hypotheses of theorem 0.2 are satisfied, then the problem

$$x'' + \frac{C'(t)}{C(t)} x' + f(t, x) = 0 \quad (0.10)$$

$$x'(0) = x(1) = 0 \quad (0.11)$$

has a solution u in E_0 such that $u(t) \leq M$.

Remarks. (a) If $\alpha(t)g(k(1-t))$ is integrable over $[0, 1]$ for all $k > 0$ then, each solution u of (0.10)–(0.11) belongs to $C^1[0, 1]$.

Proof. We remark that $(Cu')' + Cf(\cdot, u) \equiv 0$ in $(0, 1)$ and then, Cu' is uniformly continuous in $(0, 1)$. See Remark 0.3b. Thus, Cu' has a continuous extension v to $[0, 1]$ and hence, v/C is a continuous extension of u' to $[1/2, 1]$. The proof follows easily.

(b) Theorem 0.4 was considered in [3, Th. 4.1], where it was assumed that $C(t) = t^n$ and $f(t, x) = \alpha(t)x^{-p}$, for some $n \geq 1$ and $p > 0$. Note that $C(t) = t^c$ satisfies the hypotheses in Theorem 0.4, for every $c > 0$. Moreover, in this case, (0.10)–(0.11) reduces to (0.4).

(c) We first prove that Theorem 0.4 is valid, when $C(0) > 0$. Thus, the problem $[x'' + C'_n(t)x'/C_n(t) + f(t, x) = 0, x'(0) = x(1) = 0]$ has one solution u_n ; where $C_n(t) = C(1/n + (1 - 1/n)t)$. The proof will follow, from a suitable modification of the arguments in Theorem 0.1.

If (0.5) fails, we have the following alternative result:

0.5. THEOREM. Let C be as in theorem 0.4, and suppose that $f(t, x)$ is decreasing in x . If $(1-t)f(t, k(1-t))$ is integrable over $[0, 1]$ for all $k > 0$ then, (0.10)–(0.11) has a unique solution u in E_0 . Moreover, $(1-t)u'(t) \rightarrow 0$ as $t \rightarrow 1$. If in addition, $f(t, k(1-t))$ is integrable over $[0, 1]$ for all $k > 0$, then $u \in C^1[0, 1]$.

Remark. The existence part of Theorem 0.5 was proved in [4] under the additional assumptions that

- (1) $f(t, x) \rightarrow \infty$ as $x \rightarrow 0$ uniformly on compact subsets of $(0, 1)$,
- (2) $f(t, x) \rightarrow 0$ as $x \rightarrow \infty$ uniformly on compact subsets of $(0, 1)$,
- (3) f has a continuous extension to $[0, 1] \times (0, \infty)$, while the uniqueness part needs
- (4) $f(t, x)$ is strictly decreasing in x and locally Lipschitz.

To be complete, we prove a uniqueness result (Theorem 3.4 below), which contains the corresponding results in [1, 3, 4, 7]. From this corollary

we get that the problem (0.1)–(0.2) (resp. (0.1)–(0.3), (0.10)–(0.11)) has at most one solution if $f(t, x)$ is decreasing in x .

⟨1⟩ In this section we shall prove theorem 0.1. We need several notations and intermediate results.

Let J be an interval of \mathbb{R} . We denote by $C(J)$ (resp. $C^n(J)$, $n \geq 1$ integer) the space of all continuous (resp. n -times continuously differentiable) functions $u: J \rightarrow \mathbb{R}$, endowed, if J is compact, with the norm $\|u\|_0 = \sup\{|u(t)|: t \in J\}$ (resp. $\|u\|_n = \max\{\|u^{(i)}\|_0: 0 \leq i \leq n\}$). If $J = [0, 1]$, we write $C(J) = C^0$ and $C^n(J) = C^n$.

1.1. PROPOSITION. Let $u \in C^2$ satisfying the boundary conditions (0.2) and assume that there are $\alpha \in C^0$ and $g: [0, M] \rightarrow (0, \infty)$, some $M > 0$, such that $\alpha, g > 0$, g is decreasing, $0 \leq u(t) \leq M$ and

$$-u''(t) \leq \alpha(t) g(u(t)) \quad (1.1)$$

in $[0, 1]$. If $s \in (0, 1]$ denotes the unique critical point of u then

$$\int_0^{u(s)} dx/g(x) \leq \int_0^s t\alpha(t) dt. \quad (1.2)$$

Proof. See [7, Prop. 1.1].

1.2. THEOREM. Let $f_0: [0, 1] \times [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous and assume that there are α, g as in proposition 1.1 such that $0 < f_0(t, x, y) \leq \alpha(t) g(x)$ in $(0, 1) \times (0, M)$. If

$$\int_0^M dx/g(x) > \int_0^1 t\alpha(t) dt, \quad (1.3)$$

then the problem

$$x'' + f_0(t, x, x') = 0, \quad x(0) = 0, \quad \lambda x(1) + \mu x(1) = 0$$

has a solution u in C^2 such that $\|u\|_0 \leq M$.

Proof. It follows from the same arguments as in Th. 1.2 of [7].

1.3. PROPOSITION. Let $\alpha \in C(0, 1)$ be positive. If $t\alpha(t)$ is integrable over $[0, 1]$, then there is a sequence (a_n) in $(0, 1)$ such that $a_n \rightarrow 0$ and

$$a_n \int_{a_n}^1 \alpha(t) dt \rightarrow 0.$$

Proof. See [7, Prop. 1.3].

In the following, we assume that $\mu > 0$.

1.4. PROPOSITION. *Let $\gamma \in C^0$ be positive in $(0, 1)$ and for each integer $n \geq 3$, let us define $\gamma_n \in C[0, 1]$, by $\gamma_n(t) = (1 - 2/n)^2 \gamma(1/n + (1 - 2/n)t)$. If $w_n \in C^2$ is the only solution to the problem: $x'' + \gamma_n(t) = 0$, $x(0) = \lambda x(1) + \mu x'(1) = 0$, then there is $\varepsilon > 0$, such that $w_n(t) \geq \varepsilon t$ in $[0, 1]$ for $n \geq 3$.*

Proof. It follows from the same arguments as in Prop. 1.4 of [7].

Proof of Theorem 0.1. We can assume that $M < \infty$. Now, let us define $T_n(t) = 1/n + (1 - 2/n)t$; $\alpha_n(t) = (1 - 2/n)^2 \alpha(T_n(t))$, $\gamma_n(t) = (1 - 2/n)^2 \gamma(T_n(t))$, and

$$f_n(t, x) = (1 - 1/2n)^2 f(T_n(t), x + 1/n),$$

for all integers $n \geq 3$ and t in $[0, 1]$.

It is clear that

$$\gamma_n(t) \leq f_n(t, x) \leq \alpha_n(t) g(x + 1/n) \leq \alpha_n(t) g(x) \quad (1.4)$$

for $0 \leq x \leq M - 1/n$. We also have that

$$\begin{aligned} \int_0^{M-1/n} dx/g(x + 1/n) &= \int_{1/n}^M dx/g(x) \rightarrow \int_0^M dx/g(x) \\ \int_0^1 s \alpha_n(s) ds &= \int_{1/n}^{1-1/n} (s - 1/n) \alpha(s) ds \\ &< \int_{1/n}^{1-1/n} s \alpha(s) ds \rightarrow \int_0^1 s \alpha(s) ds. \end{aligned}$$

Since (0.6) holds, we can assume that

$$\int_0^{M-1/n} dx/g(x + 1/n) > \int_0^1 s \alpha_n(s) ds \quad \text{for } n \geq 3$$

and by Theorem 1.2, the problem

$$x'' + f_n(t, x) = 0, \quad x(0) = \lambda x(1) + \mu x'(1) = 0 \quad (1.5)$$

has a solution u_n such that

$$\|u_n\| \leq M - 1/n < M \quad \text{for } n \geq 3. \quad (1.6)$$

Let $t_n \in (0, 1]$ be the unique critical point of u_n . By Proposition 1.1, we get that

$$\int_0^{u_n(t_n)} dx/g(x) \leq \int_0^{T_n(t_n)} s\alpha(s) ds. \quad (1.7)$$

On the other hand, by the arguments in Thm. 1.5 of [7], we find $k > 0$ such that

$$u_n(t) \geq kt \text{ in } [0, 1] \quad \text{for } n > 2. \quad (1.8)$$

Since t_n gives the global maximum of u_n on $[0, 1]$, then (1.8) implies that $u_n(t_n) \geq u_n(1) \geq k$, for all $n \geq 3$, and from (1.7) we have $\inf\{t_n\} > 0$.

CLAIM. *Let $J = [\varepsilon, 1]$, where $0 < \varepsilon < \inf\{t_n\}$ and $\varepsilon < 1/2$. Then, there is a subsequence of $\{u_n\}$ which converges in $C^1(J)$ to a solution $w \in C^2(J)$ of (0.1). Moreover, w has a critical point.*

Proof. By (1.8) there is $\delta > 0$ such that $u_n(t) \geq \delta$ in $[\varepsilon, 1]$ and by (1.4)–(1.5),

$$-u_n''(t) \leq C\alpha_n(t) \quad \text{in } [\varepsilon, 1]$$

where $C := g(\delta)$. From this,

$$|u_n'(t) - u_n'(s)| \leq C \int_s^t \alpha_n(\tau) d\tau \leq C(1 - 2/n) \int_{T_n(s)}^{T_n(t)} \alpha(s) ds$$

for $\varepsilon \leq s < t \leq 1$; then $\{u_n'\}$ is equicontinuous in $[\varepsilon, 1]$, since α is integrable over $[\varepsilon, 1]$. Note that $T_n(\varepsilon) \geq \varepsilon$.

Moreover, for $s = t_n$ (or $t = t_n$), we get that

$$|u_n'(t)| \leq C \int_\varepsilon^1 \alpha(s) ds, \quad t \in J.$$

Thus, by Ascoli's theorem, we can assume that there exists $w \in C^1(J)$ such that $u_n \rightarrow w$ in $C^1(J)$. By (1.5) and (1.8), we have that $w \in C^2(J)$ and w is a solution to (0.1).

On the other hand, we can assume without loss of generality that $t_n \rightarrow t_0 > \varepsilon$. So, $u_n'(t_n) \rightarrow u'(t_0)$ and the proof of the claim is complete.

Now, it is not hard to prove that there is a solution u of (0.1) in $C^2(0, 1]$, which has a critical point in $(0, 1]$, such that for each compact interval J of $(0, 1]$, there exists a subsequence of $\{u_n\}$ which converges to u in $C^1(J)$. In particular, $\lambda u(1) + \mu u'(1) = 0$.

From (1.4)–(1.5), we get that

$$-u'_n(t)/g(u_n(t)) \leq \int_t^{t_n} \alpha_n(s) ds$$

for $0 < t \leq t_n$, and by integration in $[0, t] \subseteq (0, t_n]$ one has

$$\int_0^{u_n(t)} dx/g(x) \leq t \int_t^1 \alpha_n(s) ds + \int_0^t s \alpha_n(s) ds.$$

Consequently,

$$\int_0^{u(t)} dx/g(x) \leq t \int_t^1 \alpha(s) ds + \int_0^t s \alpha(s) ds,$$

and then $u(a_n) \rightarrow 0$, where $\{a_n\}$ is the sequence given by Proposition 1.3.

Let $s > 0$ be the unique critical point of u . Then, u is increasing in $(0, s)$. (Remember that $u'' < 0$) and so, $u(t) \rightarrow 0$ as $t \rightarrow 0^+$.) Thus, the proof is complete.

<2> In this section we shall prove theorem 0.4. We shall use the arguments in Corollary 1.6 of [7] to prove:

2.1. PROPOSITION. *Theorem 0.4 is true if $C(0) > 0$.*

Proof. Let us define $\psi, \phi \in C^1[0, 1] \cap C^2(0, 1)$, by

$$\psi(t) = \int_0^t C(s)^{-1} dt, \quad \phi(t) = \psi(t)/\psi(1).$$

Then, $\phi(0) = 0$, $\phi(1) = 1$, $\phi' > 0$ in $[0, 1]$, and $C\phi'' + C'\phi' \equiv 0$ in $(0, 1)$.

Now, let us define $f_0(t, x) = \phi'(\phi^{-1}(t))^{-2} f(\phi^{-1}(t), x)$ and $\alpha_0(t) = \phi'(\phi^{-1}(t))^{-2} \alpha(\phi^{-1}(t))$. Then, $f_0(t, x) \leq \alpha_0(t)g(x)$, for $0 < x < M$ and

$$\begin{aligned} \int_0^1 (1-t) \alpha_0(t) dt &= \int_0^1 (1-\phi(s)) \phi'(s)^{-1} \alpha(s) ds \\ &= \int_0^1 C(s) \left(\int_s^1 C(t)^{-1} dt \right) \alpha(s) ds \\ &\leq \int_0^1 (1-s) \alpha(s) ds, \end{aligned}$$

since $C(s) \leq C(t)$ if $s \leq t$. From this, and Theorem 0.2, the problem

$$x'' + f_0(t, x) = 0, \quad x'(0) = x(1) = 0$$

has one solution v in E_0 , such that $v(t) \leq M$. Now, it is easy to verify that $u(t) := v(\phi(t))$ is a solution to our problem.

Remark. Let $\phi \in C(0, 1)$ be positive. It is easy to prove that there is $\psi \in C[0, 1]$, such that $0 < \psi \leq \phi$ in $(0, 1)$. Thus, we can assume that the function γ in (0.5), belongs to $C[0, 1]$.

Proof of Theorem 0.4. Let us write $T_n(t) = 1/n + (1 - 1/n)t$ and $C_n(t) = C(T_n(t))$ for all integers $n > 1$ and $t \in [0, 1]$. By Proposition 2.1, the problem

$$\begin{aligned} x'' + [C'_n(t)x'/C_n(t)] + f(t, x) &= 0 \\ x'(0) &= 0, \quad x(1) = 0 \end{aligned} \quad (2.1)$$

has one solution u_n such that $u_n(t) \leq M$. Note that

$$(C_n u'_n)' + C_n f(t, u_n) = 0 \quad \text{in } (0, 1), \quad (2.2)$$

and hence

$$-C_n(\tau) u'_n(\tau) = \int_0^\tau C_n(\xi) f(\xi, u_n(\xi)) d\xi. \quad (2.3)$$

In particular, $u'_n < 0$ in $(0, 1)$ and thus, u_n is strictly decreasing in $[0, 1]$. From this,

$$-u'_n(\tau)/g(u_n(\tau)) \leq C_n(\tau)^{-1} \int_0^\tau C_n(\xi) \alpha(\xi) d\xi \quad (2.4)$$

for $0 \leq \tau \leq 1$. (Notice that $1/g(x)$ has an extension to $[0, \infty)$ which is continuous at $x = 0$.)

CLAIM 1. *There exists $k > 0$ such that $u_n(t) \geq k(1 - t)$ in $[0, 1]$, for all $n \geq 1$.*

Proof. We can assume that the function γ in (0.5) belongs to C^0 . So, for each integer $n \geq 1$, we can define

$$w_n(t) = \int_t^1 C_n(\tau)^{-1} \left(\int_0^\tau C_n(\xi) \gamma(\xi) d\xi \right) d\tau$$

It is easy to prove that w_n is the unique solution to

$$x'' + C'_n(t)x'/C_n(t) + \gamma(t) = 0, \quad x'(0) = x(1) = 0.$$

On the other hand,

$$-C_n(t) w'_n(t) = \int_0^t C_n(\xi) \gamma(\xi) d\xi \geq k_0 := \int_0^{1/2} C(\xi) \gamma(\xi) d\xi$$

in $[1/2, 1]$ and, thus, $0 < -w'_n(t) \leq 2k := k_0/C(1)$ in $[1/2, 1]$. (Note that $C_n(t) \geq C(t)$, since $T_n(t) \geq t$.) Hence, $w_n(t) \geq 2k(1-t)$ in this interval. But w_n is decreasing and so, $w_n(t) \geq w(1/2) \geq k$, in $[0, 1/2]$. From this, $w_n(t) \geq k(1-t)$ in $[0, 1]$.

Finally, $(C_n w'_n)' = -C_n \gamma \geq -C_n f(t, u_n) = (C_n u'_n)'$ and then, $C_n(t) w'_n(t) \geq C_n(t) u'_n(t)$ since $w'_n(0) = u'_n(0) = 0$. Consequently $u'_n(t) \leq w'_n(t)$, and thus $u_n(t) \geq w_n(t)$, because $w_n(1) = u_n(1) = 0$. Therefore, the proof of the claim is complete.

CLAIM 2. *For each compact interval $J := [a, b] \subseteq (0, 1)$, there exists a subsequence of $\{u_n\}$, which converges in $C^2(J)$, to a solution $w \in C^2(J)$ of (0.10).*

Proof. From (2.3) and Claim 1, we have that

$$-u'_n(t) \leq g(k(1-b)) \int_0^t \alpha(s) ds \quad \text{in } J$$

since $C_n(\xi) \leq C_n(\tau)$ if $\xi \leq \tau$; and $f(t, u_n(t)) \leq \alpha(t) g(u_n(t))$.

Thus, $\{u_n\}$ is bounded in $C^1(J)$ and by (2.1) $\{u_n\}$ is bounded in $C^2(J)$, since $\{C'_n/C_n\}$ is bounded in $C(J)$. Then, we can assume that there is v in $C^1(J)$ such that $u_n \rightarrow v$ in $C^1(J)$, and by (2.1),

$$u''_n \rightarrow -C'v'/C - f(\cdot, v) \quad \text{in } C(J).$$

The proof of Claim 2 follows easily.

Using Claim 2, we can prove the existence of a solution u to (0.10) in $C^2(0, 1)$, with the following property: For each compact interval $J \subseteq (0, 1)$, there exists a subsequence of $\{u_n\}$ which converges to u in $C^2(J)$. Therefore, u is decreasing and then, u has a continuous extension to $[0, 1]$, which we still denote by u . Note that $0 < u(t) \leq M$ in $(0, 1)$.

Let us define

$$I_1(t) = \int_t^1 (1-\xi) \alpha(\xi) d\xi, \quad I_2(t) = (1-t) \int_0^t \alpha(\xi) d\xi.$$

Integration of (2.4) in $[t, 1]$ and Fubini's theorem yield

$$\begin{aligned} \int_0^{u_n(t)} dx/g(x) &\leq \int_0^t \left(\int_t^1 C_n(\tau)^{-1} d\tau \right) C_n(\xi) \alpha(\xi) d\xi \\ &\quad + \int_t^1 \left(\int_\xi^1 C_n(\tau)^{-1} d\tau \right) C_n(\xi) \alpha(\xi) d\xi \\ &\leq I_2(t) + I_1(t), \end{aligned}$$

and then

$$\int_0^{u(t)} dx/g(x) \leq I_1(t) + I_2(t) \quad (2.5)$$

for all t in $(0, 1)$.

CLAIM 3. *There exists a sequence $\{a_n\}$ in $[1/2, 1)$ such that $a_n \rightarrow 1$ and $I_2(a_n) \rightarrow 0$.*

Proof. Assume that there exists $c > 0$ such that

$$\int_0^t \alpha(\xi) d\xi \geq c/(1-t) \quad \text{in } [1/2, 1].$$

Integrating this relation in $[1/2, r]$, for $1/2 < r < 1$, we get

$$\int_{1/2}^r (r - \xi) \alpha(\xi) d\xi \rightarrow +\infty \quad \text{as } r \rightarrow 1^-.$$

But this is a contradiction, since $(r - \xi) \alpha(\xi) \leq (1 - \xi) \alpha(\xi)$. So, the proof of Claim 3 is complete.

Let $\{a_n\}$ be as above. By (2.5) we get $u(a_n) \rightarrow 0$, and thus, $u(1) = 0$. On the other hand, by (2.3) and Claim 1, we gave that

$$0 < -u'_n(t) \leq g(k(1-t)) \int_0^t \alpha(s) ds \quad \text{if } t \in (0, 1)$$

From here, and Claim 2, $0 \leq -u'(t) \leq g(k(1-t)) \int_0^t \alpha(s) ds$ and so $u'(t) \rightarrow 0$ as $t \rightarrow 0^+$. Thus, $u \in E_0$ and the proof is complete.

<3> In this section we prove a comparison theorem, which will be basic in the proof of Theorem 0.5.

3.1. THEOREM. *Suppose that there exist $u, v \in E_0$ such that*

$$u'' + f(t, u) \geq v'' + f(t, v) \quad \text{in } (0, 1), \quad (3.1)$$

$$\lambda u(0) - \mu u'(0) \leq \lambda v(0) - \mu v'(0); \quad u(1) \leq v(1) \quad (3.2)$$

for some $\lambda, \mu \geq 0$; $\lambda + \mu > 0$. If $f(t, x)$ is decreasing in x for each fixed $t \in (0, 1)$, then $u \leq v$.

Proof. Let us define $\phi(t) = f(t, v(t)) - f(t, u(t))$ and $w = u - v$. Since (3.1) holds and $f(t, x)$ is decreasing in x , we have that

$$w'' \geq \phi, \quad \phi, w \geq 0 \quad \text{in } (0, 1), \quad (3.3)$$

and by (3.2),

$$\lambda w(0) - \mu w'(0) \leq 0; \quad w(1) \leq 0. \quad (3.4)$$

Let us fix $\tau \in [0, 1]$ such that $w(\tau) = \sup(w)$. We must prove that $w(\tau) \leq 0$. To this end, assume $w(\tau) > 0$ then, $\tau < 1$ and:

CLAIM. $w'(\tau) = 0$.

Proof. If $\tau > 0$ there is nothing to prove. Suppose now that $\tau = 0$. Then, $w'(0) \leq 0$, since $w(0) = \sup(w)$. From (3.4), $0 \leq \lambda w(0) \leq \mu w'(0) \leq 0$ and so, $\lambda = 0$ and $\mu w'(0) = 0$. Thus, the proof of the Claim is complete.

Let ξ be the point in $(\tau, 1]$ satisfying $w(\xi) = 0$ and $w > 0$ in (τ, ξ) . By (3.3), $w'' \geq 0$ in this interval and hence, the same holds for w' , since $w'(\tau) = 0$. From this, $w(\xi) \geq w(\tau) = 0$, and this contradiction ends the proof.

As an immediate consequence of Theorem 3.1, we obtain the following uniqueness result, which is related to Theorems 3 and 3.1 of [1] and [3], respectively.

3.2. COROLLARY. *If $f(t, x)$ is decreasing with respect to x , for each fixed t in $(0, 1)$, then the problem (0.1), (0.3) has at most one solution. A parallel result holds for the problem (0.1)–(0.2).*

3.3. COROLLARY. *Let $u \in E_0$ be a solution to (0.10)–(0.11) and let $v \in E_0$ be a solution to*

$$x'' + f(t, x) = 0, \quad x'(0) = x(1) = 0. \quad (3.5)$$

If $f(t, x)$ is decreasing with respect to x , and C is as in Theorem 0.4, then $u \leq v$.

Proof. From the relation $(Cu')' = Cf(t, u) = -Cf(t, u) < 0$ in $(0, 1)$, we obtain $u' < 0$ in $(0, 1)$, since $u'(0) = 0$. So, $u'' + f(u, t) \geq u'' + C'u'/C + f(t, u) = 0 = v'' + f(t, v)$, and the proof follows from Theorem 3.1.

3.4. THEOREM. *Let f, C be as in Corollary 3.3. Then, the problem (0.10)–(0.11) has at most one solution.*

Proof. Let u, v be solutions to (0.10)–(0.11) and write $w = u - v$. Then, $(Cw')'(t) = \phi(t) := C(t)[f(t, v(t)) - f(t, u(t))]; \phi \cdot w \geq 0$ and $w'(0) = w(1) = 0$.

Now, let us fix $\tau \in [0, 1]$ such that $|w(\tau)| = \sup(|w|)$ and assume $w \not\equiv 0$. Without loss of generality ($w \rightarrow -w$), we can suppose that $w(\tau) > 0$ and, as in the proof of Theorem 3.1, we get a contradiction. Thus, the proof is complete.

⟨4⟩ In this section we shall prove Theorem 0.5. We begin with the following result.

4.1. PROPOSITION. *Let $h: [0, 1] \times [0, \infty) \rightarrow (0, \infty)$ be a continuous function such that $h(t, x)$ is decreasing in x . Then, the problem*

$$x'' + h(t, x) = 0, \quad x'(0) = x(1) = 0$$

has a solution.

Proof. Let us define $H: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ by $H(t, x) = h(t, |x|)$ and notice that $0 < H(t, x) < R := 1 + \sup\{h(t, 0): t \in [0, 1]\}$.

Define $U = \{u \in C^2: \|u\|_2 < R\}$. It is easy to prove that the problem $[x'' + \lambda H(t, x) = 0, x'(0) = x(1) = 0]$ has no solution in the boundary of U for any λ in $[0, 1]$. From this, the problem $[x'' + H(t, x) = 0, x'(0) = x(1) = 0]$ has a solution u in U . Since $u'' < 0$ and $u'(0) = u(1) = 0$, then $u > 0$ in $(0, 1)$ and the proof is complete.

In the following we assume that $f(t, x)$ is decreasing in x .

4.2. THEOREM. *Assume that there exists $v, w \in E_0$ such that $w(1) = v(1) = 0$ and*

$$w''(t) + f(t, w(t)) \geq 0 \geq v''(t) + f(t, v(t)) \quad \text{in } (0, 1).$$

Assume further that there is a sequence $\{a_n\}$ in $(0, 1)$ such that $a_n \rightarrow 0$ and $w'(a_n) \geq 0 \geq v'(a_n)$ for all n . Then, the problem

$$x'' + f(t, x) = 0, \quad x'(0) = x(1) = 0 \tag{4.1}$$

has a solution $u \in E_0$.

Proof. Note that $w'(0) \geq 0 \geq v'(0)$ and, by Theorem 3.1, $w \leq v$. Now, let us choose a sequence $\{b_n\}$ in $(0, 1)$ such that $a_n < b_n$ and $b_n \rightarrow 1$, and define $T_n(t) = a_n + (b_n - a_n)t$, $w_n(t) = w(T_n(t)) - w(b_n)$, $v_n(t) = v(T_n(t)) - w(b_n)$, and $f_n(t, x) = (a_n - b_n)^2 f(T_n(t), x + w(b_n))$, for all integers $n \geq 1$ and t in $[0, 1]$.

It is clear that $w_n(1) = 0 \leq v_n(1)$, $w'_n(0) \geq 0 \geq v'_n(0)$, and

$$w''_n + f_n(t, w_n) \geq 0 \geq v''_n + f_n(t, v_n) \quad \text{in } (0, 1).$$

Let u_n be a solution to the problem

$$x'' + f_n(t, x) = 0, \quad x'(0) = x(1) = 0. \tag{4.2}$$

See Proposition 4.1. By Theorem 3.1 we have, $w_n \leq u_n \leq v_n$. In particular, $\{u_n\}$ is bounded in C^0 .

Let $J \subseteq (0, 1)$ be a compact interval, since $w_n \rightarrow w$ in $C(J)$, there exists $\varepsilon > 0$ such that $u_n(t) \geq \varepsilon$ in J for all $n > 1$ and by (4.2), $\{u_n''\}$ is bounded in $C(J)$. Consequently, $\{u_n\}$ is bounded in $C^2(J)$ and by Ascoli's Theorem, we can assume that there is z in $C^1(J)$ such that $u_n \rightarrow z$ in $C^1(J)$. By (4.2) we get that: $z \in C^2(J)$, $u_n \rightarrow z$ in $C^2(J)$ and $z'' + f(t, z) = 0$. Thus, Eq. (0.1) has one solution $u \in C^2(0, 1)$ with the following property: for each compact interval $J \subseteq (0, 1)$ there exists a subsequence of $\{u_n\}$ which converges to u in $C^2(J)$. In particular, u is bounded and decreasing and so, u has a continuous extension to $[0, 1]$, which we still denote by u . Note that $w \leq u \leq v$ and then, $u(1) = 0$.

From the relation $w_n \leq u_n \leq v_n$ we have $f_n(t, w_n) \geq f_n(t, u_n) \geq f_n(t, v_n)$ and hence $w_n'' \leq u_n'' \leq v_n''$. By integration, we have $w_n'(t) - w_n'(0) \leq u_n'(t) \leq v_n'(t) - v_n'(0)$ in $[0, 1)$ and thus,

$$w'(t) - w'(0) \leq u'(t) \leq v'(t) - v'(0) \quad \text{in } (0, 1).$$

Therefore, $u'(t) \rightarrow 0$ as $t \rightarrow 0$ and the proof is complete.

4.3. COROLLARY. *Suppose that there is $w \in E_0$ as in Theorem 4.2, and assume that $(1-t)f(t, w(t))$ is integrable over $[0, 1]$. Then, problem (4.1) has a solution.*

Proof. Since $(1-t)f(t, w(t))$ is integrable over $[0, 1]$, the problem

$$x'' + f(t, w(t)) = 0, \quad x'(0) = x(1) = 0$$

has exactly one solution v . Note that $v'' < 0$ in $(0, 1)$ and then, the same holds for v' . In particular, $v'(a_n) \leq 0$.

On the other hand, $(w-v)'' \leq 0$ and thus, $w \leq v$. Hence, $v'' + f(t, v) \leq v'' + f(t, w) = 0$; and the proof follows from Theorem 4.2.

4.4. COROLLARY. *Assume that $(1-t)f(t, k(1-t))$ is integrable over $[0, 1]$ for all $k > 0$. Then, (4.1) has a solution.*

Proof. By the arguments in Th. 0.4 of [7], there exists $\delta > 0$ such that $f(t, \delta) \geq 9\delta$ for $t \in [1/3, 2/3]$. Now, let us define $w \in C^2$ by

$$w(t) = \delta \quad \text{in } [0, 1/3]$$

$$w(t) = 27\delta(t-1/3)^4 - 18\delta(t-1/3)^3 + \delta \quad \text{in } [1/3, 2/3]$$

$$w(t) = 2\delta(1-t) \quad \text{in } [2/3, 1].$$

It is easy to prove that: $w'' \leq 0$, $w'' < 0$ in $(1/3, 2/3)$, $\min(w'') = w''(1/2) = -9\delta$, $w(1) = 0$, $w' \equiv 0$ in $[0, 1/3]$, $w' < 0$ in $(1/3, 1)$. In particular, $\max(w) = w(0) = \delta$. From this,

$$\begin{aligned} w''(t) + f(t, w(t)) &= f(t, w(t)) > 0 & \text{in } [0, 1/3] \cup [2/3, 1] \\ w''(t) + f(t, w(t)) &\geq -9\delta + f(t, \delta) \geq 0 & \text{in } [1/3, 2/3]. \end{aligned}$$

On the other hand, it is clear that $w(t) \geq k(1-t)$ for some $k > 0$ and the proof follows from Corollary 4.3, since $f(t, w(t)) \leq f(t, k(1-t))$.

Proof of Theorem 0.5. We first assume that $C(0) > 0$. With the notations of Propositions 2.1, we have

$$\begin{aligned} \min(\phi')(1-s) &\leq 1 - \phi(s) = \phi(1) - \phi(s) \leq \max(\phi')(1-s) \\ \int_0^1 (1-t)f_0(t, k(1-t)) dt &\leq \min(\phi')^{-1} \int_0^1 (1-\phi(s))f(s, k(1-\phi(s))) ds \end{aligned}$$

and hence $(1-t)f_0(t, k(1-t))$ is integrable over $[0, 1]$ for all $k > 0$. The proof, in this case, follows as in Proposition 2.1.

To prove the general case, let us define $C_n(t)$ as in theorem 0.4. Then, problem (2.1) has one solution u_n . On the other hand, let U be the unique solution to (4.1). By Corollary 3.3, $u_n \leq U$ and so, $\{u_n\}$ is bounded in C^0 .

Define now $\gamma(t) = f(t, \max(U))$. By Claim 1 of Theorem 0.4, there exists $k > 0$ such that $u_n(t) \geq k(1-t)$ in $[0, 1]$ for all $n > 1$. Thus, by the arguments in Theorem 0.4,

$$-u'_n(t) \leq \int_0^t f(s, k(1-s)) ds \quad (4.3)$$

for all $t \in [0, 1)$ and hence $\{w''_n\}$ is bounded in $C(J)$, for all compact intervals $J \subseteq (0, 1)$. From this, (0.10)–(0.11) has one solution $u \in C^2(0, 1) \cap C[0, 1]$ such that $k(1-t) \leq u(t) \leq U(t)$. In particular, $u(1) = 0$. By (4.3), $u'(t) \rightarrow 0$ as $t \rightarrow 0^+$, and so, $u \in E_0$.

On the other hand, $((1-t)Cu')' = -Cu' - (1-t)f(t, u)$, and by the arguments in Remark 0.3(b), $(1-t)C(t)u'(t)$ has a continuous extension to $[0, 1]$. Thus, the same holds for $(1-t)u'(t)$ and then $(1-t)u'(t) \rightarrow m$ for some real number $m \leq 0$. (Remember that $u'_n < 0$ in $(0, 1)$.) If $m < 0$, then there exists $c > 0$ such that $u'(t) \geq c/(1-t)$ in $(1/2, 1)$ and hence

$$\int_{1/2}^1 dt/(1-t) \leq c^{-1}u(1/2).$$

This contradiction proves that $m = 0$.

The rest of the proof follows from the arguments in Remark 0.3(b) and Remark (a) to Theorem 0.4.

REFERENCES

1. L. BOBISUD, D. O'REGAN, AND W. ROYALTY, Solvability of some nonlinear boundary value problems, *Nonlinear Anal. TMA* **12**, No. 9 (1988), 855–869.
2. A. CALLEGARI AND A. NACHMAN, Some singular differential equations arising in boundary layer theory, *J. Math. Anal. Appl.* **64** (1978), 96–105.
3. J. GATICA, V. OLIKER, AND P. WALTMAN, Singular boundary value problems for second order differential equations, *J. Differential Equations* **79** (1989), 62–78.
4. J. GATICA, G. HERNANDEZ, AND P. WALTMAN, Radially symmetric solutions of a class of singular elliptic equations, *Proc. Edinburgh Math. Soc.* **33** (1990), 169–180.
5. A. NACHMAN AND A. CALLEGARI, A nonlinear boundary value problem in theory of pseudoplastic fluids, *SIAM J. Appl. Math.* **38** (1980), 275–281.
6. S. TALIAFERRO, A nonlinear singular boundary value problem, *Nonlinear Anal. TMA* **3** (1979), 897–904.
7. A. TINEO, Existence theorems for a singular two-point Dirichlet problem, *Nonlinear Anal. TMA*, to appear.